

Model-independent determination of Z' couplings at LEP 200 *

F. DEL AGUILA

*Departamento de Física Teórica y del Cosmos, Universidad de Granada
Granada, 18071, Spain***ABSTRACT**

All couplings of a new heavy gauge boson to ordinary fermions can be determined in a model-independent way at a large e^+e^- collider for large enough statistics. No such determination is possible at LEP 200, however, because its design integrated luminosity is too low for existing and forthcoming limits on Z' interactions. At any rate, it should be possible to distinguish between models for Z' masses $< 1 \text{ TeV}$, or to set limits of this order on Z' masses for definite models. These are (preliminary) 1σ limits. We neglect systematic errors and assume an integrated luminosity of 0.5 fb^{-1} .

1. Introduction

There is no compulsory reason for expecting a new heavy Z' boson, but there is neither a reason for excluding its existence. At present there are only bounds on its effects [1-6]. The question here is: Which information on an extra gauge boson can LEP 200 provide? At a large e^+e^- collider if the Z' effects are sizeable all Z' couplings to ordinary fermions can be determined in a model-independent way [7,8]. However, the expected luminosity at LEP 200 is too low to allow for such a model-independent analysis. The typical value of the Z' mass limit, or of the discovered Z' mass, at TEVATRON when LEP 200 enters in operation is $\sim 500 \text{ GeV}$ [9]. For Z' masses $< 1 \text{ TeV}$ it is possible to distinguish between definite models at LEP 200; for specific models it is also possible to set limits on Z' masses $\sim 1 \text{ TeV}$. Our analysis is preliminary. In what follows we review present and forthcoming Z' limits (1.1) and the parametrization of Z' effects in two-fermion final state observables at e^+e^- (1.2). Our numerical estimates for Z' physics at LEP 200 are given in Section 2.

* To appear in Proceedings of the Zeuthen Workshop on Elementary Particle Theory "Physics at LEP 200 and Beyond", Teupitz, Germany, April 10-15, 1994

1.1 Present and forthcoming Z' limits

Present limits on Z' masses, $M_{Z'}$, come from the absence of an excess of $e^+e^-(\mu^+\mu^-)$ pairs with a large invariant mass at TEVATRON: *direct limits* [1,2]; and from the observation of no effect beyond the minimal standard model in precise electroweak data, mainly at LEP: *indirect limits* [3-6]. These limits for popular models, χ, ψ, η, LR , are given in Table 1 [9,10]. TEVATRON can provide limits on $M_{Z'} \sim 500(600) \text{ GeV}$ for an integrated luminosity of $25(100) \text{ pb}^{-1}$. LEP allows for a precise measurement of the Z current, providing a stringent limit on the $Z'Z_1$ mixing, $s_3 \equiv \sin\theta_3 < 0.01$ [3,6]. In models where this mixing is fixed by the Z' mass, the limit on s_3 translates into stringent $M_{Z'}$ limits: *constrained indirect limits*. When $M_{Z'}$ and s_3 are independent the Z' mass limits are modest: *unconstrained indirect limits*. The indirect limits on $M_{Z'}$ depend little on the top quark mass, m_t , except for top masses near the m_t upper bound [5].

	direct (present)	direct ($\int Ldt = 25\text{pb}^{-1}$)	direct ($\int Ldt = 100\text{pb}^{-1}$)	indirect (constrained)	indirect (unconstrained)
χ	340	470	620	670	380
ψ	320	450	600	200	200
η	340	460	610	440	210
LR	310	510	660	990	430

Table 1. Z' mass limits (in GeV).

In conclusion we will assume for our estimates at LEP 200 that a new Z' may exist with a mass $> 500 \text{ GeV}$.

1.2 Parametrization of Z' effects in two-fermion final state observables at e^+e^-

The effects of a new Z' can be observed in the two-fermion channels at LEP 200. The relevant lagrangian reads:

$$-\mathcal{L}_{NC} = e \sum_i q_i \bar{\psi}_i \gamma^\mu \psi_i A_\mu + \frac{g}{2c_W} \sum_i \bar{\psi}_i \gamma^\mu (v^i - a^i \gamma_5) \psi_i Z_{1\mu} + \frac{g}{2c_W} \sum_i \bar{\psi}_i \gamma^\mu (v'^i - a'^i \gamma_5) \psi_i Z_{2\mu}, \quad (1)$$

where the last piece describes the new Z' interaction and the first two pieces define the standard model neutral lagrangian [1]. (We assume family universality and that the $SU(2)_L$ charges T_i commute with the new charge Q' : $[Q', T_i] = 0$.) We neglect the $Z'Z_1$ mixing angle, for it is bounded, $s_3 < 0.01$, and its effects are unobservable at LEP 200. Hence, we identify Z_1 and Z_2 with the observed Z and the new Z' bosons, respectively. The three gauge bosons, γ, Z, Z' , are far off-shell for a center of mass energy of 200 GeV . The expected statistics allows for working at tree level. Radiative effects due to real Z production are important but they can be taken away with a cut on the maximum photon energy [11]. We can neglect them in a first estimate of the Z' potential of LEP 200. The main Z' effects result from the interference of the Z' amplitude,

$$\mathcal{A}^f(Z') = \frac{1}{s - M_{Z'}^2 + i\Gamma_{Z'} M_{Z'}} \left(\frac{g}{2c_W} \right)^2 \bar{u}_e \gamma^\mu (v'^e - a'^e \gamma_5) u_e \bar{u}_f \gamma_\mu (v'^f - a'^f \gamma_5) u_f, \quad (2)$$

with the γ, Z amplitudes. Neglecting the Z' width and for a fixed center of mass energy $\sqrt{s}$ ($= 200 \text{ GeV}$), the Z' amplitude and the corresponding Z' effects in $e^+e^- \rightarrow f\bar{f}$, with f any ordinary fermion, depend on five couplings [7]:

$$\epsilon_A = \frac{a'^2 e^2}{c_W^2 s_W^2} \frac{s}{M_{Z'}^2 - s}, \quad P_V^e = \frac{v'^e}{a'^e}, \quad P_L^q = \frac{v'^q + a'^q}{2a'^e}, \quad P_R^{u,d} = \frac{v'^{u,d} - a'^{u,d}}{v'^q + a'^q}. \quad (3)$$

At a large e^+e^- collider not only the pairs of charged leptons but the bottom and charm final states can be tagged. As a matter of fact, using the event shape, lepton tagging and/or vertex reconstruction LEP analyses have measured the total cross section and the forward-backward asymmetry of bottom and charm pairs [12]. Thus, the set of observables at LEP 200 includes the total cross section for charged leptons σ^l and their forward-backward asymmetry A_{FB}^l (l stands for e, μ, τ), as well as for bottom and charm quarks $R^{b,c} \equiv \frac{\sigma^{b,c}}{\sigma^l}, A_{FB}^{b,c}$, and the total cross section for hadrons $R^{had} \equiv \frac{\sigma^{had}}{\sigma^l}$. (We assume only s -channel exchange for electrons, implying $\sigma^e \sim \sigma^\mu \sim \sigma^\tau$. We do not consider tau polarization asymmetries.) If beam polarization is available, the corresponding left-right asymmetries, $A_{LR}^{l,b,c,had}, A_{LR,FB}^{l,b,c}$, double the number of observables to be measured.

σ^l and A_{FB}^l have small linear terms in P_V^e , implying that unpolarized lepton probes only determine ϵ_A and $|P_V^e|$. Whereas the polarized probes A_{LR}^l and $A_{LR,FB}^l$ are very sensitive to P_V^e , including its sign [7]. One combination of $P_L^q, P_R^{u,d}$ is fixed by R^{had} . A second combination is determined by A_{LR}^{had} if polarization is available. In any case, flavour tagging is necessary for a complete determination of the Z' couplings to quarks. $R^{b,c}$ and $A_{FB}^{b,c}$ allow for this determination even without electron beam polarization.

In Ref. [7] it is shown that a complete (model-independent) determination of the Z' couplings to ordinary fermions is possible for a Z' mass $\sim 1 \text{ TeV}$ at NLC. As we show below such a determination is not possible at LEP 200.

2. LEP 200 results

In Table 2 we gather the expected values of two-fermion final state observables at LEP 200 for different models. We assume an integrated luminosity of 0.5 fb^{-1} . Errors are only statistical. The first four columns correspond to models χ, ψ, η, LR and a Z' mass equal to 500 GeV . The fifth column gives the values of the same observables for the standard model.

	χ	ψ	η	LR	SM
$\sigma^l L_{int}$	1390 ± 40	1480 ± 40	1460 ± 40	1460 ± 40	1500 ± 40
A_{FB}^l	0.564 ± 0.020	0.542 ± 0.022	0.561 ± 0.022	0.526 ± 0.022	0.557 ± 0.021
R^{had}	7.37 ± 0.20	6.65 ± 0.19	6.86 ± 0.19	6.61 ± 0.19	6.60 ± 0.18
R^c	1.84 ± 0.06	1.78 ± 0.06	1.78 ± 0.06	1.70 ± 0.06	1.72 ± 0.06
R^b	1.23 ± 0.04	1.03 ± 0.04	1.10 ± 0.04	1.07 ± 0.04	1.05 ± 0.04
A_{FB}^c	0.657 ± 0.015	0.680 ± 0.014	0.667 ± 0.015	0.642 ± 0.015	0.670 ± 0.015
A_{FB}^b	0.585 ± 0.020	0.603 ± 0.020	0.621 ± 0.020	0.566 ± 0.020	0.604 ± 0.020
A_{LR}^l	0.038 ± 0.019	0.086 ± 0.018	0.099 ± 0.018	0.099 ± 0.018	0.086 ± 0.018
$A_{LR,FB}^l$	0.028 ± 0.019	0.064 ± 0.018	0.075 ± 0.018	0.074 ± 0.018	0.064 ± 0.018
A_{LR}^{had}	0.484 ± 0.006	0.488 ± 0.006	0.479 ± 0.006	0.561 ± 0.006	0.486 ± 0.006
A_{LR}^c	0.363 ± 0.013	0.378 ± 0.013	0.356 ± 0.013	0.453 ± 0.013	0.384 ± 0.013
A_{LR}^b	0.604 ± 0.014	0.617 ± 0.014	0.612 ± 0.014	0.677 ± 0.013	0.598 ± 0.014
$A_{LR,FB}^c$	0.187 ± 0.014	0.220 ± 0.013	0.190 ± 0.014	0.245 ± 0.014	0.219 ± 0.014
$A_{LR,FB}^b$	0.489 ± 0.015	0.533 ± 0.015	0.523 ± 0.015	0.572 ± 0.015	0.532 ± 0.015

Table 2. Values and statistical errors for two-fermion final state observables at LEP 200 ($\sqrt{s} = 200 \text{ GeV}$, $\int L dt = 0.5 \text{ fb}^{-1}$) for models χ, ψ, η, LR ($M_{Z'} = 500 \text{ GeV}$) and for the standard model.

In order to decide on the Z' potential of LEP 200 we perform a χ^2 fit to the observables in Table 2 for models χ, ψ, η, LR . l stands for e, μ, τ and counts three times. The results are given in Table 3. The errors in parentheses are for unpolarized electron beams. In both cases the errors are so large that no model-independent determination of the Z' couplings to ordinary fermions is possible.

	χ	ψ	η	LR
ϵ_A	$0.041 \pm 0.013(0.048)$	$0.069 \pm 0.043(0.076)$	$0.007 \pm 0.009(0.030)$	$0.145 \pm 0.041(0.046)$
P_V^e	$2 \pm 0.4(1.2)$	$0 \pm 0.2(10.6)$	$-3 \pm 2.2(6.7)$	$-0.148 \pm 0.1(0.4)$
P_L^q	$-0.5 \pm 0.2(0.5)$	$0.5 \pm 0.5(2.0)$	$2 \pm 1.6(5.2)$	$-0.142 \pm 0.2(0.3)$
P_R^u	$-1 \pm 0.9(1.1)$	$-1 \pm 0.7(12.5)$	$-1 \pm 0.7(1.5)$	$-6.04 \pm 7.5(13.9)$
P_R^d	$3 \pm 1.2(2.7)$	$-1 \pm 1.1(26.4)$	$0.5 \pm 0.4(2.5)$	$8.04 \pm 10.2(17.8)$

Table 3. Values of couplings in Eq. (3) and 1σ statistical errors at LEP 200 for popular models. Errors in parentheses correspond to unpolarized electron beams.

However, we can distinguish between specific models. Let us consider the class of E_6 models parametrized by β in Table 4. Performing a χ^2 fit to the observables in Table 2 for models χ, ψ, η with $M_{Z'} = 500 \text{ GeV}$, we obtain

$$\beta = 0.0 \pm 0.10(0.27), 1.57 \pm 0.10(0.28), -0.91 \pm 0.11(0.13), \quad (4)$$

respectively. Errors in parentheses correspond to unpolarized electron beams.

Figure 1:

	$\frac{g}{e}v'^i$	$\frac{g}{e}a'^i$
e	$\sqrt{\frac{2}{3}}\cos\beta$	$\frac{1}{\sqrt{6}}\cos\beta + \frac{1}{3}\sqrt{\frac{5}{2}}\sin\beta$
u	0	$-\frac{1}{\sqrt{6}}\cos\beta + \frac{1}{3}\sqrt{\frac{5}{2}}\sin\beta$
d	$-\sqrt{\frac{2}{3}}\cos\beta$	$\frac{1}{\sqrt{6}}\cos\beta + \frac{1}{3}\sqrt{\frac{5}{2}}\sin\beta$

Table 4. Vector and axial Z' couplings in Eq. (1) for a class of E_6 models. $\beta = 0, \frac{\pi}{2}, -\arctan\sqrt{\frac{5}{3}}$ for models χ, ψ, η , respectively.

For definite models we can also determine the Z' mass. A χ^2 fit to the observables in Table 2 gives

$$M_{Z'} = 500 \pm 19(25), 83(102), 43(72), 14(50), \quad (5)$$

for models χ, ψ, η, LR , respectively. In the Figure we show for the standard model observables in Table 2 the 1σ $M_{Z'}$ limits for the E_6 models in Table 4 as a function of β . The solid curve corresponds to unpolarized beams.

3. Conclusions

For LEP 200 or for any center of mass energy of the same order and an integrated luminosity $\sim 0.5 \text{ fb}^{-1}$ no model-independent determination of the Z' couplings

to ordinary fermions is possible for $M_{Z'} > 500 \text{ GeV}$. However, we can distinguish between specific models for $M_{Z'}$ fixed (and $< 1 \text{ TeV}$). Limits on $M_{Z'} \sim 1 \text{ TeV}$ can also be obtained for definite models. This analysis can be improved by including systematic errors and (initial) radiation.

Acknowledgements

We thank M. Cvetič, P. Langacker, A. Leike and S. and T. Riemann for discussions and the organizers of the workshop for their hospitality. This work was partially supported by the European Union under contract CHRX-CT92-0004 and by CICYT.

References

1. Review of Particle Properties, *Phys. Rev.* **D45** (1992) number 11, and references there in.
2. F. Abe et al., CDF Collaboration, *Phys. Rev. Lett.* **67** (1991) 2418.
3. L3 Collaboration, *Phys. Lett.* **B306** (1993) 187; ALEPH Collaboration, CERN preprint, CERN/PPE/94-30.
4. P. Langacker and M. Luo, *Phys. Rev.* **D45** (1992) 278.
5. F. del Aguila, W. Hollik, J. Moreno and M. Quirós, *Nucl. Phys.* **B372** (1992) 3; and references there in.
6. G. Altarelli et al., *Phys. Lett.* **B318** (1993) 139.
7. F. del Aguila and M. Cvetič, UPR-590-T.
8. A. Leike, DESY preprint 93-154.
9. F. del Aguila, M. Cvetič and P. Langacker, in Proceedings of the Workshop on Physics and Experiments with Linear e^+e^- Colliders, Hawaii, April 1993, Eds. F.A. Harris, S.L. Olsen, S. Pakvasa and X. Tata, p. 490; see also J.L. Hewett in the same Proceedings, p. 246.
10. See, for instance, P. Langacker and M. Luo, *Phys. Rev.* **D44** (1991) 817; F. Zwirner, *Int. J. Mod. Phys.* **A3** (1988) 49; J.L. Hewett and T.G. Rizzo, *Phys. Rep.* **183** (1989) 193; and references there in.
11. A. Djouadi, A. Leike, T. Riemann, D. Schaile and C. Verzegnassi, *Z. Phys.* **C56** (1992) 289; see also Proceedings of the Workshop on Physics and Experiments with Linear e^+e^- Colliders, Hawaii, April 1993.
12. LEP Collaborations, CERN preprint, CERN/PPE/93-157.

This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9405302v1>